

# Pinal County

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Arizona hedgehog cactus	<i>Echinocereus triglochidiatus var. arizonicus</i>	Endangered	Dark green cylindroid stem, 2.5-12 inches tall, 2-10 inches in diameter. Occurs singly or in clusters. Has 1-3 gray or pinkish central spines, the largest deflexed, and 5-11 radial spines. Flower are brilliant red along side of stem.	Gila, Pinal	3,200-5,200 ft	Ecotone between interior chaparral and madrean evergreen woodland.	Open slopes, in narrow cracks between boulders, and in understory of shrubs. Additional genetic studies have determined that the species does not occur outside of the type locality.
Bald eagle	<i>Haliaeetus leucocephalus</i>	Threatened	Large, adults have white head and tail. Height 28-38 inches; wingspan 66-96 inches. Dark with varying degrees of mottled brown plumage. Feet bare of feathers.	Gila, Graham, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	Varies	Large trees or cliffs near water (reservoirs, rivers, and streams) with abundant prey.	Some birds are nesting residents while a larger number winters along rivers and reservoirs. Once endangered (32 FR 4001, 03-11-1967; 43 FR 6233, 02-14-78) because of reproductive failures from pesticide poisoning and loss of habitat, this species was downlisted to threatened on August 11, 1995, and delisted August 8, 2007. Threatened status reinstated for Desert nesting bald eagles.
Desert pupfish	<i>Cyprinodon macularius</i>	Endangered	Small (2 inches) smoothly rounded body shape with narrow vertical bars on the sides. Breeding males blue on head and sides with yellow on tail. Females and juveniles tan to olive colored back and silvery sides.	Cochise, Graham, Maricopa, Pima, Pinal, Santa Cruz, Yavapai	< 4,000 ft	Shallow springs, small streams, and marshes. Tolerates saline and warm water.	Two subspecies are recognized: Desert Pupfish ( <i>C.m. macularis</i> ) and Quitobaquito Pupfish ( <i>C.m. eremus</i> ). Critical habitat includes Quitobaquito Springs, Pima County, portions of San Felipe Creek, Carrizo Wash, and Fish Creek Wash, Imperial County, California.
Gila chub	<i>Gila intermedia</i>	Endangered	Deep compressed body, flat head. Dark olive-gray color above, silver sides. Endemic to Gila River Basin.	Cochise, Gila, Graham, Greenlee, Pima, Pinal, Santa Cruz, Yavapai	2,000-5,500 ft	Pools, springs, cienegas, and streams.	Found on multiple private lands, including the Nature Conservancy and the Audubon Society. Also occurs on Federal and state lands and in Sonora, Mexico. Critical habitat occurs in Cochise, Gila, Graham, Greenlee, Pima, Pinal, Santa Cruz, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Lesser long-nosed bat	<i>Leptonycteris curasoae yerbabuena</i>	Endangered	Elongated muzzle, small leaf nose, and long tongue. Yellowish brown or gray above and cinnamon brown below. Tail minute and appears to be lacking. Easily disturbed.	Cochise, Gila, Graham, Greenlee, Maricopa, Pima, Pinal, Santa Cruz, Yuma	1,600-11,500 ft	Desert scrub habitat with agave and columnar cacti present as food plants.	Day roosts in caves and abandoned tunnels. Forages at night on nectar, pollen, and fruit of paniculate agaves and columnar cacti. This species is migratory and is present in Arizona usually from April to September and south of the border the remainder of the year.
Loach minnow	<i>Tiaroga cobitis</i>	Threatened	Small (<3 inches) slender, elongated fish, olive colored with dirty white spots at the base of the dorsal and caudal fins. Breeding males vivid red on mouth and base of fins.	Apache, Cochise, Gila, Graham, Greenlee, Navajo, Pinal	< 8,000 ft	Benthic species of small to large perennial streams with swift shallow water over cobble and gravel. Recurrent flooding and natural hydrograph important.	Presently found in Aravaipa Creek, Deer Creek, Turkey Creek, Blue River, Campbell Blue Creek, San Francisco River, Eagle Creek, North Fork of the East Fork Black River, and White River in Arizona, and Dry Blue Creek, Pace Creek, Frieborn Creek, the Tularosa River, West Fork Gila River, and the mainstem upper Gila River in New Mexico. Populations have been recently reintroduced in Hot Springs and Redfield canyons in Cochise and Graham counties; Fossil Creek in Gila County; and Bonita Creek in Graham County Arizona. Critical habitat (72 FR 13356-13422, March 21, 2007) found in Apache, Graham, Greenlee, and Pinal counties, Arizona, as well as portions of the Blue River, San Francisco River, Tularosa River, Negrito Creek, Pace Creek, Dry Blue Creek, Frieborn Creek, Whitewater Creek, Gila River, and its West, Middle, and East Forks in Catron, Grant, and Hidalgo counties in New Mexico.
Mexican spotted owl	<i>Strix occidentalis lucida</i>	Threatened	Medium sized with dark eyes and no ear tufts. Brownish and heavily spotted with white or beige.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai	4,100-9,000 ft	Nests in canyons and dense forests with multi-layered foliage structure.	Generally nest in older forests of mixed conifer or ponderosa pine/gambel oak type, in canyons, and use variety of habitats for foraging. Sites with cool microclimates appear to be of importance or are preferred. Critical habitat was finalized on August 31, 2004 (69 FR 53182) in Arizona in Apache, Cochise, Coconino, Gila, Graham, Greenlee, Maricopa, Navajo, Pima, Pinal, Santa Cruz, and Yavapai counties.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Mountain plover	<i>Charadrius montanus</i>	Proposed threatened	Small bird (8 – 9 ½-in), about the size of a killdeer, with longer legs and more upright posture. Light brown above with white forehead, throat, and breast but lacks the black breast band typical of many plovers. Crown may be mottled black to solid black during the breeding season and distinct black loreal stripe extending from the black bill to the eye. In winter, the crown and loreal stripe become pale brown in coloration.	Apache, Cochise, La Paz, Maricopa, Navajo, Pima, Pinal, Yuma	< 7,200 ft	Semi-desert grasslands and agricultural lands with sparse vegetation or vegetation interspersed with bare ground and flat topography.	Arizona primarily provides winter habitat for the species but some breeding occurs near Springerville.
Nichol Turk's head cactus	<i>Echinocactus horizonthalonius</i> var. <i>nicholii</i>	Endangered	Blue-green to yellowish-green, columnar, 18 inches tall, 8 inches in diameter. Spine clusters have 5 radial and 3 central spines; one curves downward and is short; 2 spines curve upward and are red or pale gray. Flowers: pink; fruit: woolly white.	Pima, Pinal	2,400-4,100 ft	Sonoran desertscrub.	Found in unshaded microsites in Sonoran desertscrub on dissected alluvial fans at the foot of limestone mountains and on inclined terraces and saddles on limestone mountain sides.
Razorback sucker	<i>Xyrauchen texanus</i>	Endangered	Large, up to 3 feet long and up to 6 lbs, high sharp-edged keel-like hump behind the head. Head flattened on top. Olive-brown above to yellowish below.	Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Pinal, Yavapai, Yuma	< 6,000 ft	Riverine and lacustrine areas, generally not in fast moving water and may use backwaters.	Big River fish also found in Horseshoe reservoir (Maricopa County). Critical habitat includes the 100-year floodplain of the river through the Grand Canyon from confluence with Paria River to Hoover Dam; Hoover Dam to Davis Dam; Parker Dam to Imperial Dam. Also Gila River from Arizona/New Mexico border to Coolidge Dam; and Salt River from Hwy 60/SR77 Bridge to Roosevelt Dam; Verde River from FS boundary to Horseshoe Lake.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Southwestern willow flycatcher	<i>Empidonax traillii extimus</i>	Endangered	Small passerine (about 6 inches) grayish-green back and wings, whitish throat, light olive-gray breast and pale yellowish belly. Two wingbars visible. Eye-ring faint or absent.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 8,500 ft	Cottonwood/willow and tamarisk vegetation communities along rivers and streams.	Migratory riparian-obligate species that occupies breeding habitat from late April to September. Distribution within its range is restricted to riparian corridors. Difficult to distinguish from other members of the Empidonax complex by sight alone. Training seminar required for those conducting flycatcher surveys. Critical habitat was finalized on October 19, 2005 (50 CFR 60886). In Arizona there are critical habitat segments in Apache, Cochise, Gila, Graham, Greenlee, Maricopa, Mohave, Pima, Pinal, and Yavapai counties.
Spikedace	<i>Meda fulgida</i>	Threatened	Small (<3 inches) slim fish with silvery sides and "spine" on dorsal fin. Breeding males are a brassy golden color.	Cochise, Gila, Graham, Greenlee, Pinal, Yavapai	< 6,000 ft	Medium to large perennial streams with moderate to swift velocity waters over cobble and gravel substrate. Recurrent flooding and natural hydrograph important to withstand invading exotic species.	Presently found in Aravaipa Creek, Eagle Creek, Verde River, and the Gila River from the San Pedro River to Ashurst-Hayden Dam in Arizona, and the Gila River and its East and West Forks in New Mexico. Populations have been recently reintroduced in Hot Springs and Redfield canyons in Cochise and Graham counties; Fossil Creek in Gila County; and Bonita Creek in Graham County Arizona. Critical habitat (72 FR 13356-13422, March 21, 2007) in Graham, Greenlee, Pinal, and Yavapai counties in Arizona, and in Catron, Grant, and Hidalgo counties in New Mexico.
Yuma clapper rail	<i>Rallus longirostris yumanensis</i>	Endangered	Water bird with long legs and short tail. Long, slender decurved bill. Mottled brown or gray on its rump. Flanks and undersides are dark gray with narrow vertical stripes producing a barring effect.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	< 4,500 ft	Fresh water and brackish marshes.	Species is associated with dense emergent riparian vegetation. Requires wet substrate (mudflat, sandbar) with dense herbaceous or woody vegetation for nesting and foraging. Channelization and marsh destruction are primary sources of habitat loss.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Acuna cactus	<i>Echinomastus erectocentrus</i> var. <i>acunensis</i>	Candidate	Less than 12 inches tall; spine clusters borne on tubercles, each with a groove on the upper surface. 2-3 central spines and 12 radial spines. Radial spines are dirty white with maroon tips. Flowers pink to purple.	Pima, Pinal	1,300-2,000 ft	Well drained knolls and gravel ridges in Sonoran desertscrub.	Immature plants distinctly different from mature plants. Immatures are disc-shaped or spherical and have no central spines until they are about 1.5 inches.
Northern Mexican Gartersnake	<i>Thamnophis eques megalops</i>	Candidate	Background color ranges from olive, olive-brown, to olive-gray. Body has three yellow or light colored stripes running down the length of the body, darker towards tail. Species distinguished from other native gartersnakes by the lateral stripes reaching the 3rd and 4th scale rows. Paired black spots extend along dorsolateral fields.	Apache, Cochise, Coconino, Gila, Graham, Navajo, Pima, Pinal, Santa Cruz, Yavapai	130-8,500 ft	Cienegas, stock tanks, large-river riparian woodlands and forests, streamside gallery forests.	Core population areas in the U.S. include mid/upper Verde River drainage, mid/lower Tonto Creek, and the San Rafael Valley and surrounding area. Status on tribal lands unknown. Distributed south into Mexico along the Sierra Madre Occidental and Mexican Plateau. Strongly associated with the presence of a native prey base including leopard frogs and native fish.
Roundtail chub	<i>Gila robusta</i>	Candidate	Member of the minnow family Cyprinidae and characterized by streamlined body shape. Color usually olive gray with silvery sides and a white belly. Breeding males develop red or orange coloration on the lower half of the cheeks and on the bases of paired fins. Individuals may reach 49.0 cm (19.3 in) but usually average 25-30 cm (9.8 - 11.8 in).	Apache, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pinal, Yavapai	1,000-7,500 ft.	Cool to warm waters of rivers and streams, often occupy the deepest pools and eddies of large streams.	Historical range of roundtail chub included both the upper and lower Colorado River basins. A 2009 status review determined that the lower Colorado River basin roundtail chub population segment (Arizona and New Mexico) qualifies as a distinct vertebrate population segment (DPS). Populations in the Little Colorado, Bill Williams, and Gila River basins are considered candidate species.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
Tucson shovel-nosed snake	<i>Chionactis occipitalis klauberi</i>	Candidate	Small snake (10-17 inches total length) in the family Colubridae, with a shovel-shaped snout and an inset lower jaw. Overall coloring mimics coral snakes, with pale yellow to cream-colored body, 21 or more black or brown saddle-like bands across the back, and orange-red saddle-like bands in between. The subspecies is distinguished from the other subspecies in that these secondary orange-red crossbands are suffused with dark pigment, making them appear brown or partly black, and the black and red crossbands do not encircle the entire body.	Maricopa, Pima, Pinal	785-1,662 ft	Sonoran Desertscrub; associated with soft, sandy soils having sparse gravel.	Found in creosote-mesquite floodplain environments, finds refuge under desert shrubs, active during crepuscular (dawn and dusk) and daylight hours.
Yellow-billed cuckoo	<i>Coccyzus americanus</i>	Candidate	Medium-sized bird with a slender, long-tailed profile, slightly down-curved bill that is blue-black with yellow on the lower half. Plumage is grayish-brown above and white below, with rufous primary flight feathers.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	< 6,500 ft	Large blocks of riparian woodlands (cottonwood, willow, or tamarisk galleries).	Neotropical migrant that winters primarily in South America and breeds primarily in the U.S. (but also in southern Canada and northern Mexico). As a migrant it is rarely detected; can occur outside of riparian areas. Cuckoos are found nesting statewide, mostly below 5,000 feet in central, western, and southeastern Arizona. Concern for cuckoos are primarily focused upon alterations to its nesting and foraging habitat. Nesting cuckoos are associated with relatively dense, wooded, streamside riparian habitat, with varying combinations of Fremont cottonwood, willow, velvet ash, Arizona walnut, mesquite, and tamarisk. Some cuckoos have also been detected nesting in velvet mesquite, netleaf hackberry, Arizona sycamore, Arizona alder, and some exotic neighborhood shade trees.

COMMON NAME	SCIENTIFIC NAME	STATUS	DESCRIPTION	COUNTY	ELEVATION	HABITAT	COMMENTS
American peregrine falcon	<i>Falco peregrinus anatum</i>	Delisted	A crow-sized falcon with slate blue-gray on the back and wings, and white on the underside; a black head with vertical "bandit's mask" pattern over the eyes; long pointed wings; and a long wailing call made during breeding. Very adept flyers and hunters, reaching diving speeds of 200 mph.	Apache, Cochise, Coconino, Gila, Graham, Greenlee, La Paz, Maricopa, Mohave, Navajo, Pima, Pinal, Santa Cruz, Yavapai, Yuma	3,500-9,000 ft	Areas with rocky, steep cliffs, primarily near water, where prey (primarily shorebirds, songbirds, and waterfowl) concentrations are high. Nests are found on ledges of cliffs, and sometimes on man-made structures such as office towers and bridge abutments.	Species recovered with over 1,650 breeding birds in the US and Canada.
Cactus ferruginous pygmy-owl	<i>Glaucidium brasilianum cactorum</i>	Delisted; petitioned for relisting	Small reddish-brown owl with a cream-colored belly streaked with reddish-brown. Males average 2.2 oz and females average 2.6 oz. Length is approximately 6.5 in., including a relatively long tail. Lacks ear tufts, and has paired black spots on the back of the head.	Pima, Pinal	< 4,000 ft	Areas of desert woodlands with tall canopy cover. Primarily found in Sonoran desert scrub and occasionally in riparian drainages and woodlands within semi-desert grassland communities. Prefers to nest in cavities in saguaro cacti but has been found in low-density suburban developments that include natural open spaces.	Not recognized as a protected taxonomic entity under the Act, but protected from direct take of individuals and nests/eggs under the Migratory Bird Treaty Act. A 2006 petition for relisting under the Act is currently being evaluated. Due to low population numbers, captive breeding research was initiated in 2006 with some success.
California brown pelican	<i>Pelecanus occidentalis californicus</i>	Delisted	Large, dark gray-brown water bird with webbed feet, pouch underneath its long bill, and wingspan of 7 ft. Adults have a white head and neck, brownish black breast, and silver gray upper parts.	Gila, La Paz, Maricopa, Mohave, Pinal, Yuma	Varies	Coastal land and islands; species found occasionally around Arizona's lakes and rivers.	Considered an uncommon transient in Arizona. Most observations recorded along the Colorado River and in the Gila Valley. Individuals known to wander up from Mexico in summer and fall. No breeding has been documented in Arizona. Delisted on December 17, 2009.